

PROJECT TITLE: Plant and Variety Selection in Spring Wheat, Winter Wheat and Barley for the Control of Foliar Diseases.

PERSONNEL: J.L.A. Eckhoff, EARC, Sidney
J.W. Bergman, EARC, Sidney
A.L. Scharen, USDA/ARS, Bozeman
D.C. Sands, MSU, Bozeman
R. McGee, MSU, Bozeman

OBJECTIVES:

1. To select germplasm of winter wheat, spring wheat and barley that has improved resistance to foliar fungal and bacterial diseases that presently occur in Montana.
2. To incorporate the resistant germplasm into winter wheat, spring wheat and barley breeding programs so that adapted varieties can be developed for release to commercial small grain growers.

SUMMARY: Foliar diseases of small grains are of economic importance to small grain producers in Montana. The most prevalent of these diseases are the fungi Septoria tritici, (Septoria leaf blotch), Septoria nodorum (Septoria glume blotch) and Pyrenophora trichostoma (tan spot), the bacteria Pseudomonas syringae and Xanthomonas translucens, and wheat streak mosaic virus. No reliable estimates are available as to the economic losses caused by these diseases, but due to their widespread occurrence, losses in aggregate are probably substantial.

Most of these foliar diseases are of particular significance under continuous cropping conditions, especially in combination with minimum tillage practices, because the disease inoculum appears to be carried over on crop residues or volunteer grain. Thus, there is an urgent need for identification of germplasm that is resistant to these diseases and development of resistant cultivars if continuous cropping and minimum tillage practices are to be applied successfully in Montana on a sustained basis.

Wheat disease nurseries have been grown at the Eastern Agricultural Research Center in cooperation with the Plant Pathology Department at Montana State University since 1975. Each year, 250-700 spring wheat lines and 150-500 winter wheat lines have been evaluated for disease resistance potential. Materials for the nursery are provided by Dr. A. Scharen of the Pathology Department and come from various environments, including Israel, Lebanon, Tunisia, Kenya, India, Japan, USSR, Yugoslavia, Algeria, Spain, Mexico, Brazil, Chile, Argentina, Australia and other wheat producing countries. All the disease evaluations are made by personnel from the Plant Pathology Department at Montana State University. In addition to disease ratings, other agronomic characteristics of these wheat varieties are rated by the staff of the Eastern Agricultural Research Center. The most promising lines are harvested each year and are used as a source of resistance in the breeding programs at Montana State University.

One hundred fifty-nine winter wheat lines were planted under irrigated conditions during the 1989-90 growing year. The winter wheat lines were read for disease reaction on June 19, 1990. Winter survival was good. Septoria infection was moderate, and readings of resistance were possible.

Two hundred eleven spring wheat lines were planted under irrigated conditions in 1990. The spring wheat lines were read for disease on July 5, 1990. Septoria infection was moderate, and readings of resistance were possible.

A barley disease resistant recurrent selection population has been grown at the Eastern Agricultural Research Center since 1979 in cooperation with the Plant Pathology Department at Montana State University. The population was provided by Dr. David Sands. Each year, Dr. Harold Bockelman evaluated disease reaction at Sidney and selected resistant plants. The population was primarily selected for resistance to Xanthomonas translucens (bacterial leaf streak), although selection was for resistance to any disease organism. Each year, several thousand plants were grown and evaluated for disease resistance. Resistant plants were harvested and the seed produced were planted in a winter nursery in Arizona. Plants in Arizona were intercrossed and the resulting seed were planted back in Sidney in the following spring as the disease resistant recurrent selection population.

Lines from the disease resistant population were selected for good agronomic characters. Eight hundred plant selections from the barley disease resistant recurrent selection population were planted under dryland conditions in 1990. Rows were evaluated for agronomic characteristics and the best rows were harvested for seed increase.

RESULTS: Septoria tritici(leaf blotch) and Septoria nodorum(glume blotch) were observed in the irrigated winter wheat and spring wheat nurseries, and lines were evaluated for resistance to these diseases. Severity of these diseases was moderate in both winter wheat and spring wheat. No other diseases were noted.

FUTURE PLANS: Because Dr. Al Scharen will retire soon, the Septoria winter and spring wheat nurseries will be discontinued in 1991.

Spring barley advanced lines which were selected from the disease resistant recurrent selection population and increased in Sidney last year will be grown in preliminary yield trials in 1991. In addition, recurrent selection will continue in the parent population and lines with disease resistance and good agronomic characters will be selected for seed increase and yield testing.