

PROJECT TITLE: Winter and Spring Wheat Variety Performance Evaluation Under Northern Montana Conditions on the Basis of Gross Production Value as Influenced by Yield, Protein, and Market.

PROJECT LEADER: Gregg R. Carlson, Agronomist - Havre

PROJECT PERSONNEL: Jim Christianson, Executive VP, MWBC - Great Falls
 Cheryl Tuck, Information Officer, MWBC - Great Falls
 P.L. Bruckner, Breeder/Geneticist (WW) - Bozeman
 L.E. Talbert, Breeder/Geneticist (SW) - Bozeman
 J.E. Berg, Research Associate (WW) - Bozeman
 S.P. Lanning, Research Associate (SW) - Bozeman
 Cooperating County Extension Agents
 Cooperating Landowners

OBJECTIVES:

It is the objective of this project to bring quality and quantity together to allow the forces of market value to influence evaluation of winter and spring wheat varieties under varying cropping conditions in northern Montana.

RESULTS:

Average annual PNW quotes for Hard Red Winter wheat at 10, 11, 12, and 13 percent protein for the 10-year period 1991-2000 are graphed in Figure 1. The PNW annual market averages for the same period for Dark Northern Spring wheat at 13, 14, and 15 percent protein (plus values for 12 percent protein first available in 1994), are graphed in Figure 2. Both graphs include values along the top axis reflecting the average annual \$/bu price spread between the minimum and maximum protein levels for which quotes are consistently given.

'Gross Dollar Return' comparisons are graphically presented in Figures 3. through 7. reflecting research plot response data for the locations and years below:

Figure No.	Cropping System & Crop	Research Location	No. of Varieties Included	No. of Data Years Included	No. of Calendar Years	Calendar Years Spanned
3	Flw-WW	NARC-Havre	9	10	11	1990-2000
*3a	Flw-WW	NARC-Havre	7	7	7	1994-2000
4	Flw-SW	NARC-Havre	6	10	11	1990-2000
*4a	Flw-SW	NARC-Havre	7	6	6	1995-2000
5	Flw-SW	Turner	7	10	11	1990-2000
*5a	Flw-SW	Turner	8	8	9	1992-2000
*5b	Flw-SW	Turner	8	6	7	1994-2000
6	Flw-SW	Loring	8	5	5	1996-2000
7	Flw-SW	Loma	9	3	3	1998-2000

Analyses span the maximum number of calendar years, up to 10, for which data exists for a specific location and variety set. Figures marked with an asterisk (*) denote those for which a reduced number of data years were used in the analyses for purposes of including new or otherwise popular variety releases having fewer data years available. In contrast to the 'Comparable Average' method of comparing varietal performance graphs in this report reflect only analyses where each variety shown was actually grown at that particular location during all years listed. Thus, values included are actual in terms of agronomic yield and associated gross return based on protein content and average annual market performance for each year.

It must be kept in mind that in addition to the influence of price variability; crop fertility, plant available water, and

numerous other factors can dramatically affect gross dollar return. All trials included were fertilized. No attempt has been made here to consider fertilizer or other input costs and their subsequent effect on 'net' return. Plant available water estimates and soil fertility data are available for these studies. Climatic and nursery management data details for each off-station trial featured here are included with the associated agronomic evaluations in a separate report. Most Havre data represents a minimum 'applied' nitrogen rate of 70 lbs actual N/ac. It should be further understood that management plays a key role in affecting dollar return - be it associated with enhanced product quality, quantity or shrewd marketing skills.

SUMMARY:

Producers are well aware of the impact protein premiums can have on overall market value, but are troubled by the fact that the market has generally not been consistent in terms of rewarding growers for producing high quality wheat. The potential for discount associated with low quality has likely had more bearing on production management than have positive incentives in the form of premiums for quality above average. In the past decade, average annual premiums for 10-13 percent protein winter wheat and 12-15 percent protein spring wheat have varied from as little as 1.7 cents to as much as 49.3 cents per point increase per bushel. Producers have encouraged researchers to evaluate potential new practices in terms of dollars and cents. Such is never easy; and this particular effort toward quantifying wheat variety performance on the basis of total dollar return was no exception.

Working with MWBC, the Research Center initiated development of a 'Gross Dollar Return' database in 1988 utilizing a limited approach involving Wednesday markets only. By 1989, daily market spreadsheets were made available by MWBC with some file development assistance for previous years provided by NARC. At present, full market data for the years 1973-2000 have been made available to the Research Center.

For each research location a multi-year, average gross market value per acre was determined for each selected variety. Such values were based on gross return for actual yield at the lowest consistently quoted protein level plus added gross return for protein premium, if any. The sum of the two values then represents the gross return per acre in a given market year. Calculations were made for each year the varieties were under evaluation at a particular location. The values were then tested via simple analysis of variance with data years as replications.

It should be noted that the current procedure affords no mechanism for appropriate adjustment of gross return where protein content is either below that termed as "minimum quoted" (10 percent for winter wheat and 13 percent for spring wheat, 12 percent for spring wheat beginning with 1994) or above that termed as "maximum quoted" (13 percent for winter wheat and 15 percent for spring wheat). Thus, discounts for protein below the minimum quoted - or added premiums sometimes available for protein above normal quote maximums, cannot be reflected in these data. Due to fertilization, situations where protein levels were below minimum are extremely rare in these research databases. However, situations where protein exceeded the maximum level for which market quotes were available are common in these data. Thus, in cases where proteins for 'average protein performing' varieties in a particular trial are at the maximum level for which a market quote exists; entries with higher protein are not benefited by additional premium as they may have been in a commercial marketing situation.

One must also remain aware that the marketing periods chosen for these analyses can have pronounced effects on the results due to obvious year differences in overall market price and premium spreads. Not unlike most crop evaluation procedures, but perhaps even more important in this case, data reliability increases with additional years of observation. At present, it would appear that a minimum of four to five years should be involved for meaningful comparison via this system.

In 1994, Carlson initiated a new "paired" trial series at Turner whereby 16-23 varieties each of wheat and barley were evaluated for five years under both low and optimum nitrogen fertility. Abridged results of that 5-year study in terms of agronomics and fertilizer economics are posted at Northern Agricultural Research Center's website in the agronomy research section at <http://agadsrv.msu.montana.edu/narc/agronomy.htm>

FUTURE PLANS:

The Research Center plans to continue work with MWBC and wheat breeders in further developing and refining the use of these data with agricultural producers. Regression or other more sophisticated means of analysis could be

introduced in work with these data. Use of additional data sets representing conditions of lower fertility will also be important to refine the assessment of economic benefits associated with production of high quality varieties.

We have considered an entire revamping of our market matrices to reflect a more logical market year than is represented by the current CY arrangement. Very little new production year wheat is marketed in northern Montana until at least mid-August. Thus it may be more logical to associate a years' agronomic data with 12 months following harvest - perhaps beginning September 1. Such would be a bit frustrating, as agronomic data for a year could not be economically compared until nearly a year later. However, such would more accurately represent the real world. If we take this approach, we would now be able to analyze only up through the 1999 crop. We wouldn't be incorporating the 2000 crop data into the system until summer 2001. Furthermore, it may be sensible to assign value weighting to individual months within the annual average on the basis of traditional market volume during those months.

We will continue to explore ways in which to improve the use of actual market data in the comparison of wheat varieties and production practices.

Average Annual Market Quotes

* (\$/Bu - Hard Red Winter Wheat)

Pacific Northwest Delivery




Figure 1.

MSU/AES/NARC-Carlson

* Average of All Market Days/Market Year
** \$/Bu Difference Between 10 & 13% Protein

Average Annual Market Quotes

* (\$/Bu - Dark Northern Spring Wheat)

Pacific Northwest Delivery




Figure 2.

MSU/AES/NARC-Carlson

* Average of All Market Days/Market Year
** \$/Bu Difference Between 13 & 15% Protein

Gross Return - Fallow Winter Wheat Northern Ag Research Center - Havre, Montana ¹10-Yr Means (1990-2000)




Figure 3.

MSU/AES/NARC-Havre
Ref=00-3512 P=.0000 CV2=2.82%

¹1992 Nursery Lost to Hail
LSD (P<=.05), Gross Return = \$17.28 / ac
Prices = PNW Average Annual Market/Year

Gross Return - Fallow Winter Wheat Northern Ag Research Center - Havre, Montana 7-Yr Means (1994-2000)




Figure 3a.

MSU/AES/NARC-Havre
Ref=00-3522 P=.0039 CV2=3.11%

LSD (P<=.05), Gross Return = \$20.98 / ac
Prices = PNW Average Annual Market/Year

Gross Return - Fallow Spring Wheat Northern Ag Research Center - Havre, Montana ¹10Yr Means (1990-2000)




Figure 4.

MSU/AES/NARC-Havre

Ref=00-3112 P=.0001 CV2=2.89%

¹1992 Nursery Lost to Hail

LSD (P<=.05), Gross Return = \$ 16.78 / ac

Prices = PNW Average Annual Market/Year

Min.Quote@12%Pro (1994-00), 13% (1990-93)

Gross Return - Fallow Spring Wheat Northern Ag Research Center - Havre, Montana 6-Yr Means (1995-2000)




Figure 4a.

MSU/AES/NARC-Havre

Ref=00-3122 P=.0003 CV2=3.42%

LSD (P<=.05), Gross Return = \$21.20 / ac

Prices = PNW Average Annual Market/Year

Gross Return - Fallow Spring Wheat Leon Cederberg Farm – Turner, Montana ¹10-Yr Means (1990-2000)




Figure 5.

MSU/AES/NARC-Havre
Ref00-9911 P=.0000 CV2=2.40%

¹Not Planted in 1999 (too wet until too late)
LSD (P<=.05), Gross Return = \$ 13.00 / ac
Prices = PNW Average Annual Market/Year

Gross Return - Fallow Spring Wheat Leon Cederberg Farm – Turner, Montana ¹8-Yr Means (1992-2000)




Figure 5a.

MSU/AES/NARC-Havre
Ref=00-9921 P=.0000 CV2=2.51%

¹Not Planted in 1999 (too wet until too late)
LSD (P<=.05), Gross Return = \$ 14.87 / ac
Prices = PNW Average Annual Market/Year

Gross Return - Fallow Spring Wheat Leon Cederberg Farm – Turner, Montana 16-Yr Means (1994-2000)




Figure 5b.

MSU/AES/NARC-Havre
Ref00-9931 P=.0012 CV2=2.94%

¹Not Planted in 1999 (too wet until too late)
LSD (P<=.05), Gross Return = \$ 17.09 / ac
Prices = PNW Average Annual Market/Year

Gross Return - Fallow Spring Wheat Flansaas/Lumsden Farm - Loring, Montana 5-Yr Means (1996-2000)




Figure 6.

MSU/AES/NARC-Havre
Ref=00-9905 P=.0001 CV2=1.81%

LSD (P<=.05), Gross Return = \$ 8.37 / ac
Prices = PNW Average Annual Market/Year

Gross Return - Fallow Spring Wheat

McKeever Farm & Seed, Inc. – Loma, Montana

3-Yr Means (1998-2000)




Figure 7.

MSU/AES/NARC-Havre

Ref=00-9907 P=.0752 CV2=4.40%

LSD (P<=.05), Gross Return = \$ 17.33 / ac
Prices = PNW Average Annual Market/Year